



HIP SURVEILLANCE PROGRAM

for Children with Cerebral Palsy

British Columbia Consensus Statement on Hip Surveillance for Children with Cerebral Palsy

Information for Health Care Professionals
Caring for Children with Cerebral Palsy

LEAD BENEFACTOR



SUMMARY

The British Columbia Consensus Statement on Hip Surveillance for Children with Cerebral Palsy was established by a group of over 60 multidisciplinary professionals from all regions of the province. The group included parents of children with cerebral palsy (CP), pediatric orthopaedic surgeons, physiotherapists, occupational therapists, developmental pediatricians, pediatricians, family physicians, nurses, a radiologist and radiographer, policy makers and health administrators. Meetings were held in May 2011 and January 2012. Attendees established consensus regarding the desire to create a BC hip surveillance program for children with CP and reached consensus on the commencement, frequency, and discharge criteria for surveillance using a voting system. When required, discussion continued until consensus could be reached. This booklet describes the established consensus.

AUTHORS OF THIS REPORT: BC Hip Surveillance Planning Committee (in alphabetical order)

Janice Duivesteyn, Program Manager, Sunny Hill Health Centre for Children
Nancy Lanphear, Senior Medical Director, Sunny Hill Health Centre for Children
Mary Lou Matthews, Provincial Lead, Network Operations, Child Health BC
Tanja Mayson, Physiotherapist, Sunny Hill Health Centre for Children
Stacey Miller, Physiotherapist, BC Children's Hospital
Kishore Mulpuri, Orthopaedic Surgeon, BC Children's Hospital
Maureen O'Donnell, Executive Director, Child Health BC
Lori Roxborough, Therapy Department Director, Sunny Hill Health Centre for Children
Suzanne Steenburgh, Program Manager, BC Children's Hospital
Esias van Rensburg, Developmental Pediatrician, Sunny Hill Health Centre

CITING THIS REPORT

BC Hip Surveillance Planning Committee. British Columbia Consensus Statement on Hip Surveillance for Children with Cerebral Palsy: Information for health care professionals caring for children with cerebral palsy. Child Health BC, Vancouver, BC. 2012.

www.childhealthbc.ca/hips.

ACKNOWLEDGEMENTS

The authors would like to thank all of the meeting participants who contributed to the creation of the BC Consensus Statement on Hip Surveillance for Children with Cerebral Palsy and an implementation model. Participants are listed in meeting reports that are available on the Child Health BC website (www.childhealthbc.ca/hips).

The authors also wish to acknowledge the work of colleagues from Australia and Sweden which was used as a basis for this Consensus Statement. We thank Professor Kerr Graham, Pam Thomason, Kate Willoughby, Meredith Wynter, and Kelly Kerr for sharing their invaluable knowledge and experience.

TABLE OF CONTENTS

Introduction and Objective.....	1
Evidence for Hip Surveillance	1
Inclusion	1
Hip Displacement	2
Level of Risk	2
Motor Function	2
Gait Pattern	4
Assessments	4
Clinical Examination	5
Radiological Assessment	5
Recommended Frequency of Hip Surveillance	7
GMFCS I & II	7
Hemiplegia Group IV	7
GMFCS Level III	8
GMFCS IV & V	8
Referral to Orthopaedic Surgeon	9
Hip Surveillance Discharge Criteria	9
How To Use This Consensus Statement.....	9
Recommended Frequency of Hip Surveillance - Quick Guide	10
References	11
Appendix A – Abbreviations	13

British Columbia Consensus Statement on Hip Surveillance for Children with Cerebral Palsy

INTRODUCTION AND OBJECTIVE

Hip surveillance is the process of identifying and monitoring critical early indicators of progressive hip displacement.¹ Hip displacement, or subluxation, is the gradual movement of the femoral head laterally from under the acetabulum. A hip is dislocated when the femoral head is completely displaced from under the acetabulum.

Children with cerebral palsy (CP) are at risk for hip displacement. The objective of this Consensus Statement is to outline recommendations for hip surveillance to ensure that children with CP receive appropriate screening and are referred to a pediatric orthopaedic surgeon at the appropriate time to minimize or prevent complications associated with hip dislocations. Hip displacement is often silent, with no physical signs or symptoms. Left untreated, displaced or dislocated hips may cause pain, decreased hip range of motion, decreased sitting, standing, or walking tolerance, and difficulty with personal care. Timely orthopaedic management is critical to those children identified through surveillance as having progressive displacement. This document does not address the orthopaedic management of progressive hip displacement. The intervention should be tailored to the needs of the individual child.

Surveillance practices in Australia and Sweden^{1,2}, best available research evidence, and expert opinion were used to establish this Consensus Statement. This document was created for health care professionals caring for children at risk for hip displacement.

EVIDENCE FOR HIP SURVEILLANCE

Early identification of hip displacement through surveillance has been demonstrated to be an effective means of reducing the incidence of hip dislocation in children with CP. A systematic review on the evidence for hip surveillance found surveillance is an effective means of identifying hip displacement.³ Hip surveillance programs, in combination with timely orthopaedic management, have been shown to decrease or prevent the incidence of hip dislocations in children with CP.⁴⁻⁶

INCLUSION

All children with CP should be included in a hip surveillance program. CP is defined as:

“a group of permanent disorders of the development of movement and posture, causing activity limitation, that are attributed to non-progressive disturbances that occurred in the developing fetal or infant brain. The motor disorders of cerebral palsy are often accompanied by disturbances of sensation, perception, cognition, communication, and behaviour, by epilepsy, and by secondary musculoskeletal problems”.^{7(p. 9)}

CP is not defined by the underlying cause of the condition. All non-progressive disturbances of the fetal or infant brain occurring in the pre-natal, peri-natal and post-natal period, up to the age of 2 years, can lead to CP. For example, children with a genetic anomaly, a chromosomal abnormality, a metabolic condition, or an acquired brain injury resulting from meningitis, encephalitis, or a stroke in early life can also be diagnosed with CP if they have the motor findings described in the above definition. In these children, we simply understand why they have CP.

Disorders of the spinal nerves (i.e. spina bifida), peripheral nerves (i.e. spinal muscular atrophy), muscles (i.e. muscular dystrophy), or mechanical origins (i.e. arthrogryposis) are not considered CP and, therefore, this Consensus Statement is not applicable.

HIP DISPLACEMENT

The hip is a ball and socket joint, with the acetabulum making up the 'socket' and the femoral head making up the 'ball' (Figure 1a). In a healthy hip the femoral head rests completely in the acetabulum. Hip displacement, also called subluxation, refers to the gradual movement of the femoral head laterally from under the acetabulum (Figure 1b). A hip is dislocated when the femoral head is completely displaced from under the acetabulum (Figure 1c). Delayed or absent weight bearing, limitations in gross motor function, and abnormal muscle forces around the hip joint affect the development of the proximal femur and hip joint.




Figure 1a: Normal Hip




Figure 1b: Displaced Hip




Figure 1c: Dislocated Hip

LEVEL OF RISK

Hip displacement is the second most common deformity in children with CP.⁸ The overall incidence of hip displacement in children with CP has been found to be approximately one-third (26-35%).^{2,9-11}

Motor Function

Hip displacement has been shown to be directly related to a child's gross motor function, as described by the child's Gross Motor Function Classification System (GMFCS) level, and is, therefore, used as a basis for this Consensus Statement (Figure 2).^{2,9-11}




Figure 2: Incidence of hip displacement (Migration Percentage >30%) by GMFCS Level⁹

The GMFCS is a five level classification system for children with CP that is based on self initiated movement.^{12,13} It was originally created in 1997 and was expanded and revised in 2007.^{12,13} The GMFCS-Expanded & Revised is available for download, free of charge, at https://www.canchild.ca/system/tenon/assets/attachments/000/000/058/original/GMFCS-ER_English.pdf.

Classifying a child's GMFCS level requires familiarity with the child and their usual performance of motor skills but no formal training is required. It can be completed in only a few minutes. Distinctions between levels are based on functional limitations, the need for hand held mobility devices or wheeled mobility, and, to a lesser extent, quality of movement. Each level has been given a title that reflects the method of mobility typical for children at that level after 6 years of age:

- GMFCS I: Walks without Limitations
- GMFCS II: Walks with Limitations
- GMFCS III: Walks Using a Hand-Held Mobility Device
- GMFCS IV: Self-Mobility with Limitations; May Use Powered Mobility
- GMFCS V: Transported in a Manual Wheelchair

Separate age bands exist as classification of motor function is dependent upon age. These age bands include:

- before 2nd birthday,
- between 2nd and 4th birthday,
- between 4th and 6th birthday,
- between 6th and 12th birthday, and
- between 12th and 18th birthday.

Children under the age of 2, if born premature, should be classified based on their corrected age. Expectations for gross motor function differ by age so it is important to consult the User Instructions each time a child's motor function is classified.


Gait Pattern

Most children with hemiplegia are classified at GMFCS level I or II. As such, they are at low risk for hip displacement and discharged from surveillance prior to skeletal maturity. However, children with a Winters, Gage and Hicks (WGH) Group IV hemiplegic gait must be followed until skeletal maturity (from now on referred to as Group IV gait).

Winters, Gage and Hicks¹⁴ described the classification of hemiplegic gait into four gait patterns (Figure 3). Group IV gait involves more marked proximal involvement of the hip. The WGH classification system is only based on changes in the sagittal plane but many children with hemiplegia also have deviations in the coronal and transverse plane. It is these children who have changes in all three planes of motion that should be included in hip surveillance. For the purposes of hip surveillance, Group IV gait pattern is characterized by:

- a flexed hip and an anterior pelvic tilt (sagittal plane),
- an adducted hip (coronal plane), and
- hip internal rotation (transverse plane).¹⁴

Children with this gait pattern are at risk of progressive hip displacement that typically occurs later than children with bilateral CP.¹ Risk of hip displacement in children with bilateral cerebral palsy is not related to their gait pattern. Their level of risk is determined by GMFCS level only.



Reprinted with permission of IOS Press and the Royal Children's Hospital, Melbourne, Australia¹⁵

Figure 3: Winters, Gage, and Hicks Hemiplegia Gait Patterns.^{14, 15}

ASSESSMENTS

Hip surveillance requires both clinical and radiological review. Clinical examination is an important component of hip surveillance but hip displacement cannot be based on clinical assessment alone.^{4,16}

The frequency of the clinical and radiological examinations is dependent upon a child's risk for hip displacement and is, therefore, determined by the child's GMFCS level and, for children with hemiplegia, the child's gait pattern. The recommended frequency of clinical and radiological examinations is described on pages 7 and 8.

Clinical Examination^a

This Clinical Exam is solely for the purpose of hip surveillance and is to be completed by the child's physiotherapist. If a child does not have a physiotherapist, it is to be completed by a designated health care professional familiar with the assessments.

Classify:

- Determine GMFCS level
- Identify Group IV hemiplegia

Measure: (before measuring, please see the Clinical Exam Instructions, available at www.childhealthbc.ca/hips, for the standardized method of completing the measures involved).

- Hip abduction range of motion measured with hips at 0° flexion and knees fully extended (R2 value)
- Dynamic contracture of adductors using the Modified Tardieu Scale (R1 value)
- Modified Thomas test for hip flexion contracture

Ask the Child's Parent or Primary Caregiver:


- Does your child have hip pain? You may notice this when changing your child's position, when you move your child's leg, or when looking after your child's personal care.
- Do you have more difficulty looking after your child during activities that involve moving the hip, such as personal care, dressing, bathing, etc?
- Has there been a decrease in your child's ability to walk, sit, or stand, which is related to the hip?
- Who is your family physician/pediatrician?

Radiological Assessment

The radiological measure used to monitor hip displacement is migration percentage (MP). MP is defined as the percentage of the ossified femoral head outside of the lateral margin of the ossified acetabulum (Figure 4).¹⁷ An antero-posterior (AP) radiograph of the pelvis taken in a standardized position is required to accurately measure the migration percentage (Figure 5).¹⁷⁻¹⁹ MP is affected by the amount of abduction or adduction of the leg so the leg should be positioned in neutral. Measurement of the MP requires that the triradiate cartilages be visible and therefore anterior and posterior pelvic tilt must be corrected.


Reimers reported that the 90th percentile for MP in typically developing children at 4 years of age was 10%.¹⁷ A hip is considered 'at risk' when the MP is greater than 30%. Evidence supports measurement of MP by a single, experienced examiner.^{20,21}

^a The Clinical Exam component of the Consensus Statement does not replace the need for regular, comprehensive musculoskeletal assessment as a component of overall management.



Reproduced with permission and copyright © of the British Editorial Society of Bone and Joint Surgery⁴.

Figure 4: Measurement of Migration Percentage⁴




Reproduced with permission and copyright © of the British Editorial Society of Bone and Joint Surgery⁴.

Figure 5: Standardized positioning for antero-posterior radiograph

RECOMMENDED FREQUENCY OF HIP SURVEILLANCE


GMFCS I & II^{13,22}




- At each Clinical Exam, verify GMFCS level and identify children with a Group IV gait
- If GMFCS level has changed or child is identified as having a Group IV gait, ongoing surveillance according to confirmed classification
- Initial Clinical Exam at identification of CP
- Review annually with Clinical Exam
- Review at 5 years of age with Clinical Exam and AP pelvic radiograph
- If radiograph findings are normal at 5 years, discharge from surveillance

Hemiplegia Group IV^{14, 15}

- Until 5 years of age, surveillance as per recommendations for children at GMFCS I & II
- After 5 years of age, Clinical Exam and AP pelvic x-ray 12 monthly until skeletal maturity




GMFCS Level III^{13, 22}




- At each Clinical Exam, verify GMFCS level; if GMFCS level has changed, ongoing surveillance according to confirmed classification
- Initial Clinical Exam at identification of CP
- Clinical Exam and initial AP pelvic radiograph at 24 months of age
- Clinical Exam and AP pelvic radiograph 12 monthly until 6 years of age
- After 6 years of age, until skeletal maturity, review with:
 - Clinical Exam 12 monthly
 - AP pelvic radiograph 24 monthly

GMFCS IV & V^{13, 22}



- At each Clinical Exam, verify GMFCS level; if GMFCS level has changed, ongoing surveillance according to confirmed classification
- Initial Clinical Exam at identification of CP
- Clinical Exam and initial AP radiograph at 24 months of age
- Clinical Exam and AP pelvic radiograph 6 monthly until 6 years of age
- After 6 years of age, continue Clinical Exam and AP pelvic radiograph 12 monthly until skeletal maturity



REFERRAL TO ORTHOPAEDIC SURGEON

The intention of this Consensus Statement is that review by a pediatric orthopaedic surgeon occurs at the appropriate time. A referral to an orthopaedic surgeon should occur in the following situations:

- Migration Percentage > 30%
- End of range hip abduction (R2) $\leq 30^\circ$ when measured with hips at 0° flexion and knees extended
- Deterioration or asymmetry in:
 - Hip abduction (R1 or R2)
 - Modified Thomas test
- Positive answer to any one of the three questions in the Clinical Exam
- Any other clinical concern that is felt to be related to the hip

An aim of hip surveillance is that orthopaedic review occurs at the appropriate time when treatment options are available. Every child referred to orthopaedic services should be managed with an individualized management plan, which may or may not include ongoing hip surveillance.¹ Children who have surgery for hip displacement or dislocation should return to surveillance post operatively until reaching skeletal maturity.

HIP SURVEILLANCE DISCHARGE CRITERIA

Hip displacement can occur during the pubertal growth spurt and thus children at risk must be followed until skeletal maturity.⁹ The closure of the triradiate cartilage on the AP radiograph is used as the prime indicator of skeletal maturity in the BC Consensus Statement.¹




Children that are at GMFCS levels I & II, excluding those with a Group IV gait pattern, are at low risk for hip displacement, and are followed until the age of 5 years.

HOW TO USE THIS CONSENSUS STATEMENT

This consensus statement is for use by individuals caring for children with CP, within the Child Health BC Hip Surveillance Program. It is intended for education and to assist in clinical decision making. **Individual clinicians are to use their own clinical judgment in decision making about individual clients.** Should you have questions, contact the child's physician or orthopaedic surgeon.

British Columbia Consensus on Hip Surveillance for Children with Cerebral Palsy¹

QUICK GUIDE

Classification	Age in Years										Continue Until Bones Stop Growing (on X-ray)	
	ID	2	2.5	3	3.5	4	4.5	5	5.5	6		
 <p>GMFCS I^{2,3} GMFCS II^{2,3}</p>								 				
 <p>GMFCS III^{2,3}</p>		 		 		 		 		 	 	Every year Every 2 years
 <p>GMFCS IV^{2,3} GMFCS V^{2,3}</p>		 	 	 	 	 	 	 	 	 	 	Every year Every year
 <p>Group IV Hemiplegic Gait^{4,5}</p>								 		 	 	Every year Every year

Legend: GMFCS: Gross Motor Function Classification System²
ID: Identification/Diagnosis of cerebral palsy
Group IV Hemiplegic Gait: Child walks with one hip turned and pulled inward⁴



Clinical Exam



X-Ray

References:

1. BC Hip Surveillance Planning Committee. British Columbia Consensus Statement on Hip Surveillance for Children with Cerebral Palsy. Child Health BC, Vancouver, BC, Canada. 2012.
2. Palisano R et al. Content validity of the expanded and revised Gross Motor Function Classification System. *Dev Med Child Neurol.* 2008;50:744-50.
3. Illustrations Version 2. Reid B, Willoughby K, Harvey A, Graham HK. The Royal Children's Hospital, Melbourne, Australia.
4. Winters TF, Gage JR, Hicks R. Gait patterns in spastic hemiplegia in children and young adults. *J Bone Joint Surg (Am).* 1987;69:437-441.
5. Illustrations printed with permission of IOS Press and The Royal Children's Hospital, Melbourne, Australia.

REFERENCES

1. Wynter M, Gibson N, Kentish M, Love S, Thomason P, Graham HK. The consensus statement on hip surveillance for children with cerebral palsy: Australian standards of care. *J Pediatr Rehabil Med*. 2011;4(3):183-95.
2. Hagglund G, Lauge-Pedersen H, Wagner P. Characteristics of children with hip displacement in cerebral palsy. *BMC Musculoskelet Disord*. 2007;8:101-106
3. Gordon GS, Simkiss DE. A systematic review of the evidence for hip surveillance in children with cerebral palsy. *J Bone Joint Surg [Br]*. 2006;88-B: 1492-1496.
4. Dobson F, Boyd RN, Parrott J, Natrass GR, Graham HK. Hip surveillance in children with cerebral palsy. Impact on the surgical management of spastic disease. *J Bone Joint Surg [Br]*. 2002;84-B:720-726.
5. Elkamil AI, Andersen GL, Hägglund G, Lamvik T, Skranes J, Vik T. Prevalence of hip dislocation among children with cerebral palsy in regions with and without a surveillance programme: a cross sectional study in Sweden and Norway. *BMC Musculoskelet Disord*. 2011;12:284-290.
6. Hagglund G, Andersson S, Duppe H, Lauge-Pedersen H, Nordmark E, & Westbom L. Prevention of dislocation of the hip in children with cerebral palsy. The first ten years of a population-based prevention programme. *J Bone Joint Surg [Br]*. 2005;87-B:95-101.
7. Rosenbaum P, Paneth N, Leviton A, Goldstein M, Bax M. A report: The definition and classification of cerebral palsy April 2006. *Dev Med Child Neurol*. 2007;49:8-14.
8. Cornell MS. The hip in children with cerebral palsy: Predicting the outcome of soft tissue surgery. *Clin Orthop*. 1995;340:165-171.
9. Soo B, Howard JJ, Boyd RN, Reid SM, Lanigan A, Wolfe R, Reddihough D, Graham HK. Hip displacement in cerebral palsy. *J Bone Joint Surg [Am]*. 2006;88:121-129.
10. Connelly A, Flett P, Graham HK, Oates J. Hip surveillance in Tasmanian children with cerebral palsy. *J Pediatr Child Health*. 2009;45:437-443.
11. Kentish M, Wynter M, Snape N, Boyd R. Five year outcome of state-wide hip surveillance of children and adolescents with cerebral palsy. *J Pediatr Rehabil Med*. 2011;4:205-217.
12. Palisano R, Rosenbaum P, Walter S, Russell D, Wood E, Galuppi B. Development and reliability of a system to classify gross motor function in children with cerebral palsy. *Dev Med Child Neurol*. 1997;39:214-223.
13. Palisano RJ, Rosenbaum P, Bartlett D, Livingston MH. Content validity of the expanded and revised Gross Motor Function Classification System. *Dev Med Child Neurol*. 2008;50:744-750.
14. Winters T, Gage J, Hicks R. Gait patterns in spastic hemiplegia in children and adults. *J Bone Joint Surg [Am]*. 1987;69:437-441.
15. Reprinted from *Journal of Pediatric Rehabilitation Medicine. An Interdisciplinary Approach 4*. Wynter M, Gibson N, Kentish M, Love S, Thomason P, Graham HK. The consensus statement on hip surveillance for children with cerebral palsy: Australian standards of care, 183-95, 2011, with permission from IOS Press and The Royal Children's Hospital, Melbourne, Australia.
16. Scrutton D, Baird G. Surveillance measures of the hips of children with bilateral cerebral palsy. *Arch Diseases Child*. 1997;76:381-384.
17. Reimers J. The stability of the hip in children: a radiological study of results of muscle surgery in cerebral palsy. *Acta Orthop Scand*. 1980;184:1-100.
18. Hagglund G, Lauge-Pedersen H, Persson M. Radiographic threshold values for hip screening in cerebral palsy. *J Child Ortho*. 2007;1:43-47
19. Scrutton D. The early management of hips in cerebral palsy. *Dev Med Child Neurol*. 1989;31:108-116.
20. Parrott J, Boyd RN, Dobson F, Lancaster A, Love S, Oates J, Wolfe R, Natrass GR, Graham HK. Hip displacement in spastic cerebral palsy: repeatability of radiological measurement. *J Pediatr Orthop*. 2002;22:660-667.
21. Faraj S, Atherton WG, Stott NS. Inter- and intra-measurer error in the measurement of Reimers' hip migration percentage. *J Bone Joint Surg [Br]*. 2004;86-B:434-7.
22. Illustrations Version 2 by Reid B, Willoughby K, Harvey A, Graham K. The Royal Children's Hospital, Melbourne, Australia.

Appendix A

ABBREVIATIONS

AP	antero-posterior
BC	British Columbia
CP	cerebral palsy
GMFCS	Gross Motor Function Classification System
MP	migration percentage
WGH	Winter, Gage, and Hicks



For more information on the Child Health BC Hip Surveillance Program, visit www.childhealthbc.ca/hips.

To speak with the Hip Surveillance Coordinator, call 604-875-2345 or 1-888-300-3088, extension 4099, or email hips@cw.bc.ca.

March 2016